

Improving snow compaction and *firn densification* on E3SM's ice sheets

*Schneider, A. M.*¹, *Zender, C. S.*¹, & *Price, S. F.*²

¹UC Irvine, Department of Earth System Science, Irvine, CA 92697

²Fluid Dynamics and Solid Mechanics Group, LANL, Los Alamos, NM 87545

1. Introduction

NEWS AND VIEWS · 30 SEPTEMBER 2020

The worst is yet to come for the Greenland ice sheet

An assessment of past, present and future ice loss from the Greenland ice sheet shows that rates of loss in the twenty-first century will be much higher than those at any time during the past 11,700 years.

Andy Aschwanden 

Andy Aschwanden is at the Geophysical Institute, the University of Alaska Fairbanks, Fairbanks, Alaska 99775, USA.

 Contact

Search for this author in:

[Pub Med](#)

[Nature.com](#)

[Google Scholar](#)

By **Jenessa Duncombe**  10 December 2019

The interior of Greenland's ice sheet doesn't usually make headlines: It's a layer of compact snow and glacial ice at high elevations that typically doesn't contribute to runoff that drives sea level rise.


But a new study suggests that this may change: More runoff may come from Greenland's interior because of a newly discovered phenomenon called ice slabs. The slabs are layers of ice that exist just below the snow's surface, where porous snow usually sits. Ice slabs can extend for tens of kilometers and grow to over 16 meters thick.

Ice slabs can extend for tens of kilometers and grow to over 16 meters thick.

Left: Briner et al., featured in a recent edition of *Nature* (<https://doi.org/10.1038/d41586-020-02700-y>), predict unprecedented Greenland Ice Sheet (GrIS) mass loss this century, contributing 2.4-9.9 cm to sea level rise. GrIS mass loss is dominated by its change in surface mass balance (SMB).


Left: *EOS* article covering observations of “ice slabs” forming near the GrIS surface (<https://eos.org/articles/a-new-source-of-sea-level-rise-from-greenland-ice-slabs>). It is unclear how *firn densification* will impact future SMB.

2. E3SM Land Model (ELM) development




Left: ELM 1D-snowpack grid: default in v2 (top, max. of 5 layers) and with “*use_extrasnowlayers*” (bottom, max. of 16 layers). The max. number of snow layers and the max. allowed snowpack depth have been increased to improve the simulation of snow compaction and *firn densification* needed for future *SMB* studies.

3. GrIS firn density profiles




0 10 20 30 40 50 60
E3SM land snow water equivalent (m)

Above: Map of GrIS ELM snowpack and SUMup (Montgomery et al., 2018) locations.



Left: Variation of density in depth, including density measurements (SUMup dataset, scattered), as simulated with the CESM Land Model v5 snowpack configuration (CLM-12snl, light-blue), and with our expanded E3SM land snowpack configuration (ELM-16snl, dark-green/black). SUMup density measurements are sorted into 12 groups, each representing the nearest ELM node (indicated by subplot lat-lon coordinates) for ease of geographical comparisons.


4. Polar firn densification through 100 years



Left: Variation of densities (top) and advective strain rates ($\times 10^{-10}$, bottom) as a function of firn age in CLM-12snl, (light-blue), and in (ELM-16snl, dark-green/black) and compared against the model of Herron & Langway, 1980 (see colorbar label). Results are sorted into 3 groups representing the mean annual temperature indicated in subplot titles.

5. Conclusions and future work

- Developments enable deep (~50 m) firn simulations
- Top 10 m firn densities compare well with measurements
- Optimization (in “ELM-16snl”) improves simulation of deeper (>10 m) *firn densification*



Left: 2 years of snow accumulation in v2 with “*use_extrasnowlayers*”. Ongoing simulations are testing and initializing the snowpack and firn conditions needed to calculate GrIS *SMB* in E3SM.

6. Acknowledgements

- This work is supported by the U.S. Department of Energy's Scientific Discovery Through Advanced Computing (LANL-520117) and Earth System Model Development (DE-SC0019278) programs and used resources of the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility.
- Model source code was obtained from the Energy Exascale Earth System Model (E3SM) project, sponsored by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research.
- We thank all the scientists, software engineers, and administrators who contributed to the development of E3SM.
- We also thank those working on the Community Earth System Model and netCDF software developed by UCAR/Unidata.

7. References

- Briner, J.P., Cuzzone, J.K., Badgeley, J.A. et al. Rate of mass loss from the Greenland Ice Sheet will exceed Holocene values this century. *Nature* 586, 70–74 (2020).
<https://doi.org/10.1038/s41586-020-2742-6>
- Herron, M., & Langway, C. (1980). Firn Densification: An Empirical Model. *Journal of Glaciology*, 25(93), 373-385. doi:10.3189/S0022143000015239
- Montgomery, L., Koenig, L., and Alexander, P.: The SUMup dataset: compiled measurements of surface mass balance components over ice sheets and sea ice with analysis over Greenland, *Earth Syst. Sci. Data*, 10, 1959–1985, <https://doi.org/10.5194/essd-10-1959-2018>, 2018.
- van Kampenhout, L., Lenaerts, J. T. M., Lipscomb, W. H., Sacks, W. J., Lawrence, D. M., Slater, A. G., & van den Broeke, M. R. (2017). Improving the representation of polar snow and firn in the Community Earth System Model. *Journal of Advances in Modeling Earth Systems*, 9, 2583– 2600.
<https://doi.org/10.1002/2017MS000988>